

ON THE DISTANCE BETWEEN PERFECT NUMBERS.

PH. ELLIA

ABSTRACT. If n, m are distinct perfect numbers and if $|n - m| = \delta$ it is known that $\delta > 1$. We show that $\delta \nmid n$. Then we prove that there exist infinitely many triangular numbers $\delta \equiv 3 \pmod{12}$ such that $|x - y| = \delta$ has no solutions with both x and y perfect numbers.

1. INTRODUCTION

An integer $n \in \mathbb{N}$ is said to be perfect if $\sigma(n) = 2n$ where σ is the sum of divisors function. By results of Euler every even perfect number has the form $n = 2^{p-1}(2^p - 1)$ where $2^p - 1$ is prime, whereas every odd perfect number is of the form $n = q^{4b+1} \cdot \prod p_i^{2a_i}$, q, p_i distinct primes, $q \equiv 1 \pmod{4}$; in particular if n is an odd perfect number, then $n \equiv 1 \pmod{4}$. It is still unknown if odd perfect numbers exist (for some recent results see for example [4], [5]).

In their very stimulating paper [3], Luca and Pomerance show that if the *abc*-conjecture is true, then the equation $x - y = \delta$ has a finite number of solutions with x, y perfect numbers when δ is odd or if x and y are both squarefull.

An obvious remark is in order here: if one could prove that the distance between two perfect numbers is always even, then it would follow that any perfect number is even.

It has also been proved that if n and m are perfect, then $|n - m| > 1$ (cf [2]).

Here we show that if n and $n + \delta$, $\delta > 1$, are both perfect, then $\delta \nmid n$, see Proposition 4.

It is known that modular properties imply that an integer $\delta \equiv \pm 1 \pmod{12}$ can't be the distance between two perfect numbers (Remark 6). It is natural to ask if there are other infinite series of numbers δ such that the equation $|x - y| = \delta$ has no solutions with x, y both perfect.

We show (Theorem 8) that there exist infinitely many triangular numbers $\delta \equiv 3 \pmod{12}$ with this property.

2. A PROPERTY OF THE DISTANCE BETWEEN TWO PERFECT NUMBERS.

We first recall the following ([2]):

Lemma 1. *If n and m are perfect numbers and if $\delta := |n - m|$, then $\delta > 1$.*

2010 *Mathematics Subject Classification.* 11A99.

Key words and phrases. Perfect numbers, distance, triangular numbers.

We have:

Lemma 2. *Let n and $n + \delta$ be perfect numbers. If $\delta \geq 2$ is even, then $\delta \nmid n$.*

Proof. This is clear if n is odd. So assume $n = 2^{p-1}(2^p - 1)$, $n + \delta = 2^{q-1}(2^q - 1)$, where $P = 2^p - 1$ and $Q = 2^q - 1$ are primes and $q > p$. Then $n + \delta > 2n$, which implies that $\delta > n$, so $\delta \nmid n$. \square

Lemma 3. *Let n , $n + \delta$, be perfect numbers. If $|\delta| > 1$ is odd, then $\delta \nmid n$.*

Proof. We may assume that n is even. So $n = 2^{p-1} \cdot Q$, where $Q = 2^p - 1$ is prime. Then $|\delta| = Q$ and so $n + \delta = Q(2^{p-1} \pm 1)$. But by Euler, we must have $Q^2 \mid n + \delta$, a contradiction. \square

Gathering everything together, we get:

Proposition 4. *If n, m are distinct perfect numbers and if $\delta := |n - m|$, then $\delta \nmid n$.*

Remark 5. *If n, m are perfect numbers, it is easy to show that $\delta := |n - m| > 2$. Moreover if $\delta \equiv 3 \pmod{4}$ and if $n > 6$, it can be proved that $(\delta, n) = 1$.*

3. SPECIAL VALUES OF THE DISTANCE BETWEEN TWO PERFECT NUMBERS.

As stated in the introduction, using known modular results we have:

Remark 6. *An integer $\delta \equiv \pm 1 \pmod{12}$ can't be the distance between two perfect numbers.*

Indeed this follows from the fact that any even perfect number is $\equiv 6$ or $4 \pmod{12}$ (Euler's theorem), while any odd perfect number is $\equiv 1$ or $9 \pmod{12}$ by Touchard's theorem ([6], [1]).

We introduce some notations:

In the sequel we denote by $\delta = b(b - 1)/2$ a triangular number such that:

$b = 4m + 2 = 48s + 46$ ($m = 12s + 11$) and $s \not\equiv 2 \pmod{3}$.

Observe that $\delta = (24s + 23)(48s + 45) \equiv 3 \pmod{12}$.

Lemma 7. *Keep notations as above. If n and $n + \delta$ are perfect numbers, then n is odd and:*

$$n = q^{4\beta+1} \cdot \prod p_u^{2\alpha_u} = (2^{p-1} - 2m - 1) \cdot (2^p + 4m + 1)$$

where $Q = 2^p - 1$ is prime, $q \equiv 1 \pmod{4}$ and where q and the p_u 's are distinct odd primes.

Moreover if $l = 8m + 3$ is prime then:

$$(1) \quad 2^{p-1} - 2m - 1 = \prod p_i^{2a_i} =: u^2$$

$$(2) \quad 2^p + 4m + 1 = q^{4\beta+1} \cdot \prod p_j^{2a_j} =: qv^2$$

and $q \equiv 5 \pmod{8}$.

Proof. First let us show that n is odd and $n + \delta$ is even. Since $\delta \equiv 3 \pmod{4}$, this is clear if $n > 6$. If $n = 6$, $6 + \delta$ is odd. Since $\delta \equiv 0 \pmod{3}$, $3^2 \mid 6 + \delta$, but this is impossible since $s \not\equiv 2 \pmod{3}$.

So $n + \delta = Q(Q + 1)/2$ where $Q = 2^p - 1$ is prime and $n = q^{4\beta+1} \cdot \prod p_u^{2\alpha_u}$. We have:

$$\frac{Q(Q+1)}{2} - \frac{b(b-1)}{2} = \frac{(Q+b)(Q-b+1)}{2}$$

It follows that $n = A \cdot B$ with $A = (2^{p-1} - 2m - 1)$, $B = (2^p + 4m + 1)$. Since $B - 2A = 8m + 3 = l$, the g.c.d., G , of A, B is 1 or l since l is prime by assumption.

If $G = l$, since $l \equiv 3 \pmod{4}$, $l \neq q$. If $q \nmid B$, then $B = 2^p + 4m + 1 = l^c \cdot \prod p_j^{2a_j}$ with $c \in \{1, 2a-1\}$ ($l^{2a} \parallel n$). Since, by assumption, $m \equiv 3 \pmod{4}$, $2^p + 4m + 1 \equiv 5 \pmod{8}$. Since $l \equiv 3 \pmod{8}$ and c is odd, we get a contradiction. We conclude that $q \mid B$, hence $B = l^c \cdot q^{4\beta+1} \cdot \prod p_j^{2a_j}$ and we get: $5 \equiv B \equiv 3 \cdot q^{4\beta+1} \equiv 3 \cdot q \pmod{8}$. Since $q \equiv 1 \pmod{4}$, this is impossible. This shows that $G = 1$.

So one of A, B is a square and the other is divisible by $q^{4\beta+1}$. Assume $B = 2^p + 4m + 1 = \prod p_j^{2a_j}$, then $5 \equiv 2^p + 4m + 1 \equiv \prod p_j^{2a_j} \equiv 1 \pmod{8}$, so A is a square, $q \mid B$ and $q \equiv 5 \pmod{8}$. \square

Now we have:

Theorem 8. *There exist infinitely many triangular numbers δ , with $\delta \equiv 3 \pmod{12}$, such that the equation $x - y = \delta$ has no solution with x and y both perfect numbers.*

Proof. We keep notations and assumptions of Lemma 7. We have $l = 8m + 3 = 96s + 91$. If $s = 3a$, $l = 288a + 91$; if $s = 3a + 1$, $l = 288a + 187$. Since $(288, 91) = 1$ and $(288, 187) = 1$, by Dirichlet's theorem there are infinitely many values of s for which l is prime.

If l is prime, by Lemma 7: $2^{p-1} - 2m - 1 = x^2$ and $2^p + 4m + 1 = qy^2$. Adding these two equations we get:

$$3 \cdot 2^{p-1} + 2m = x^2 + qy^2$$

it follows, since $m \equiv 2 \pmod{3}$, that:

$$1 \equiv x^2 + qy^2 \pmod{3}$$

- If $x^2 \equiv 0 \pmod{3}$: then $2^{p-1} - 2m - 1 \equiv 0 \pmod{3}$ and it follows that $2^{p-1} \equiv 2 \pmod{3}$, which is impossible because $p - 1$ is even.

- If $y^2 \equiv 0 \pmod{3}$: then $2^p + 4m + 1 \equiv 0 \pmod{3}$, which implies $2^p \equiv 0 \pmod{3}$: impossible.
- We conclude that $x^2 \equiv y^2 \equiv 1 \pmod{3}$, but then we get $q \equiv 0 \pmod{3}$, i.e. since q is prime $q = 3$, which is also impossible because $q \equiv 1 \pmod{4}$

We conclude that if $\delta = (24s + 23) \cdot (48s + 45)$ and if $l = 96s + 91$ is prime, then $|x - y| = \delta$ has no solutions with both x and y perfect numbers. \square

Remark 9. *Of course we can get other examples. If $b = 30, m = 7$, then $8m + 3 = 59$ is prime. Since $\delta = 435$, n is odd and Lemma 7 applies. Hence it suffices to show that $2^{p-1} - 15$ is not a square. Now if $2^{p-1} - 15 = x^2$, we have $(2^{\frac{p-1}{2}} - x)(2^{\frac{p-1}{2}} + x) = 15 = 3 \cdot 5$ and it follows that $p = 5$ or $p = 7$. Since these values of p are easily excluded in our context, we conclude that $\delta = 435$ can't be the distance between two perfect numbers.*

The question of whether an odd triangular number can be the distance between two perfect numbers remains open.

REFERENCES

- [1] Holdener, J.A.: *A theorem of Touchard on the form of odd perfect numbers*, The American Mathematical Monthly, Vol. 109, No. 7, 661-663 (2002)
- [2] Luca, F.-Coglan, F.B.-: *Problems and solutions: Can two consecutive numbers both be perfect?* The American Mathematical Monthly, Vol. 108, No. 1, 80-81 (2001)
- [3] Luca, F.-Pomerance, C.: *On the radical of a perfect number*, New York J. Math. **16**, 23-30 (2010)
- [4] Nielsen Pace, P.: *Odd perfect numbers have at least nine distinct prime factors*, Math. Comp., **76**, 2109-2126 (2007)
- [5] Ochem, P.-Rao, M.: *Odd perfect numbers are greater than 10^{1500}* , Math. Comp., 81, 1869-1877 (2012)
- [6] Touchard, J.: *On prime numbers and perfect numbers*, Scripta Mathematica, 19, 35-39 (1953)

DIPARTIMENTO DI MATEMATICA, 35 VIA MACHIAVELLI, 44100 FERRARA
E-mail address: phe@unife.it