

**ON THE COHOMOLOGY OF RANK TWO VECTOR BUNDLES ON \mathbb{P}^2
AND A THEOREM OF CHIANTINI AND VALABREGA.**

PH. ELLIA

ABSTRACT. We show that a normalized rank two vector bundle, E , on \mathbb{P}^2 splits if and only if $h^1(E(-1)) = 0$. Using this fact we give another proof of a theorem of Chiantini and Valabrega. Finally we describe the normalized bundles with $h^1(E(-1)) \leq 4$.

1. INTRODUCTION.

We work over an algebraically closed field of characteristic zero. It follows from a famous theorem of Horrocks ([9]) that a rank two vector bundle E on \mathbb{P}^n , $n \geq 2$, splits if and only if $H_*^i(E) := \bigoplus_{k \in \mathbb{Z}} H^i(E(k)) = 0$, for $0 < i < n$. This has been improved: as a consequence of another famous theorem by Evans-Griffith, under the same assumptions, E splits if and only if $H_*^1(E) = 0$ (see [5]). Along these lines, on \mathbb{P}^3 , there is a remarkable result:

Theorem 1. (Chiantini-Valabrega [3])

Let \mathcal{F} be a rank two vector bundle on \mathbb{P}^3 .

- (1) If $c_1(\mathcal{F}) = 0$, then \mathcal{F} splits if and only if $h^1(\mathcal{F}(-1)) = 0$.
- (2) If $c_1(\mathcal{F}) = -1$, then \mathcal{F} splits if and only if $h^1(\mathcal{F}(-1)) = 0$ or $h^1(\mathcal{F}) = 0$ or $h^1(\mathcal{F}(1)) = 0$.

It is natural to ask if there is a similar result on \mathbb{P}^2 and indeed there is: let E be a normalized (i.e. $-1 \leq c_1(E) \leq 0$) rank two vector bundle on \mathbb{P}^2 , then E splits if and only if $h^1(E(-1)) = 0$. Furthermore this is the best possible result. Indeed if $E = \Omega(1)$, then $h^1(E(m)) = 0, \forall m \neq -1$, but E is indecomposable. Actually this result follows from a more general fact: *with notations as above, we have $h^1(E(k)) \leq h^1(E(-1)), \forall k \in \mathbb{Z}$* (see Theorem 2). The proof of Theorem 2 is quite easy using standard vector bundles techniques. This statement has certainly been (unconsciously) known since a long time but, as far as I know, hasn't been put in evidence. That's a pity because it has some interesting consequences. For example we show how to recover Theorem 1 from it. (For another application see [6].)

In the last section, after some general considerations, we describe rank two vector bundles on \mathbb{P}^2 with $h^1(E(-1)) \leq 4$.

2. VARIATIONS ON A THEOREM OF CHIANTINI AND VALABREGA.

Let us take some notations and recall some basic facts.

Date: January 4, 2019.

2010 Mathematics Subject Classification. 14F05.

Key words and phrases. Rank two vector bundles, projective plane, cohomology.

If F is a rank two vector bundle on \mathbb{P}^n , $n \geq 2$, then $c_1(F(m)) = c_1(F) + 2m$ and $c_2(F(m)) = c_1(F)m + c_2(F) + m^2$. A rank two vector bundle E is *normalized* if $-1 \leq c_1(E) \leq 0$. In this case we will denote by c_1, c_2 its Chern classes.

In the sequel E will always denote a normalized rank two vector bundle with Chern classes c_1, c_2 .

The integer r_E (or just r if no confusion can arise) is defined as follows $r = \min\{k \in \mathbb{Z} \mid h^0(E(k)) \neq 0\}$. In other words r is the least twist of E having a section. Let $s \in H^0(E(r))$. If s does not vanish, then $E \simeq \mathcal{O}(-r) \oplus \mathcal{O}(r + c_1)$. If s vanishes, by minimality, its zero locus $(s)_0 = Z$, has codimension two and we have an exact sequence: $0 \rightarrow \mathcal{O} \rightarrow E(r) \rightarrow \mathcal{I}_Z(2r + c_1) \rightarrow 0$. The subscheme Z is l.c.i. and $\deg(Z) = c_2(E(r))$.

The bundle E is said to be *stable* if $r > 0$. If $r \leq 0$ we will say that E is *not stable* (it can be semi-stable if $c_1 = 0$).

If E is not stable and indecomposable, then $h^0(E(r)) = 1$, hence Z is uniquely defined.

Finally we recall Riemann-Roch theorem: If F is a rank two vector bundle on \mathbb{P}^2 with Chern classes c_1, c_2 , then

$$\chi(F) = 2 + \frac{c_1(c_1 + 3)}{2} - c_2.$$

In particular if E is a normalized rank two vector bundle on \mathbb{P}^2 with Chern classes c_i , then:

$$(1) \quad \chi(E(k)) = \frac{c_1}{2}(c_1 + 2k + 3) + (k + 1)(k + 2) - c_2.$$

If \mathcal{F} is a rank two normalized vector bundle on \mathbb{P}^3 with Chern classes c_i , then:

$$(2) \quad \begin{aligned} \text{If } c_1 = 0 : \quad & \chi(\mathcal{F}(k)) = -c_2(k + 2) + \frac{1}{3}(k + 1)(k + 2)(k + 3) \\ \text{If } c_1 = -1 : \quad & \chi(\mathcal{F}(k)) = \frac{1}{6}(k + 1)(k + 2)(2k + 3) - \frac{c_2}{2}(2k + 3) \end{aligned}$$

Now we can prove the main result of this section:

Theorem 2. *Let E be a rank two normalized vector bundle on \mathbb{P}^2 . Then:*

- (1) $h^1(E(k)) \leq h^1(E(-1)), \forall k \in \mathbb{Z}$.
- (2) E splits if and only if $h^1(E(-1)) = 0$.

Proof. (1) We may assume E indecomposable. If E is not stable we have an exact sequence: $0 \rightarrow \mathcal{O} \rightarrow E(r) \rightarrow \mathcal{I}_Z(2r + c_1) \rightarrow 0$, with $r \leq 0$ and $Z \subset \mathbb{P}^2$ a non-empty zero-dimensional subscheme. Twisting by $\mathcal{O}(-r - 1)$ and taking cohomology we get: $h^1(E(-1)) = h^1(\mathcal{I}_Z(r - 1 + c_1))$. Since $r - 1 + c_1 < 0$, $h^1(\mathcal{I}_Z(r - 1 + c_1)) = h^0(\mathcal{O}_Z) =: \deg(Z)$. Now for any k , the exact sequence above shows that $h^1(E(k)) \leq h^1(\mathcal{I}_Z(k + r + c_1))$. Since $h^1(\mathcal{I}_Z(m)) \leq h^0(\mathcal{O}_Z), \forall m$ (consider $0 \rightarrow \mathcal{I}_Z(m) \rightarrow \mathcal{O}(m) \rightarrow \mathcal{O}_Z \rightarrow 0$), we are done.

Now assume E is stable. Let $L \subset \mathbb{P}^2$ be a general line and consider the exact sequence $0 \rightarrow E(m - 1) \rightarrow E(m) \rightarrow E_L(m) \rightarrow 0$. Since $E_L = \mathcal{O}_L \oplus \mathcal{O}_L(c_1)$ (Grauert-M'ulich theorem, see [9]), if $m \leq -1$, $h^0(E_L(m)) = 0$ and $h^1(E(m - 1)) \leq h^1(E(m))$. It follows that $h^1(E(m)) \leq h^1(E(-1))$ if $m \leq -1$. If $m \geq 0$, by Serre duality $h^1(E(m)) = h^1(E^*(-m - 3)) = h^1(E(-m - 3 - c_1))$ and again $h^1(E(m)) \leq h^1(E(-1))$.

(2) Of course (2) follows from (1) and Horrocks' theorem, but let us give a simpler argument. If E is not stable arguing as in (1), we get $\deg Z = 0$, hence $Z = \emptyset$ and E splits. It remains to show that $h^1(E(-1)) > 0$ if E is stable. By stability $\chi(E(-1)) = -h^1(E(-1))$. By Riemann-Roch, if $h^1(E(-1)) = 0$, we get $c_2 = 0$. Now $\chi(E) = 2$ if $c_1 = 0$ (resp. 1 if $c_1 = -1$). It follows that $h^2(E) > 0$. But $h^2(E) = h^0(E^*(-3)) = h^0(E(-c_1 - 3)) = 0$, by stability. Hence $h^1(E(-1)) \neq 0$. \square

Remark 3. *This is the best possible result in the sense that for any $m \neq -1$, there exists an indecomposable rank two vector bundle, E , with $h^1(E(m)) = 0$: just take $E = \Omega(1)$.*

Remark 4. *Let's consider an unstable rank two vector bundle, E , with $c_1(E) = -1$. Arguing as above we see that $h^1(E) = 0$ implies that E splits.*

Assume now $h^1(E(1)) = 0$. We have $0 \rightarrow \mathcal{O} \rightarrow E(r) \rightarrow \mathcal{I}_Z(2r - 1) \rightarrow 0$. Twisting by $\mathcal{O}(-r + 1)$ we get: $0 \rightarrow \mathcal{O}(-r + 1) \rightarrow E(1) \rightarrow \mathcal{I}_Z(r) \rightarrow 0$ it follows that $h^1(\mathcal{I}_Z(r)) = 0$. Now consider $0 \rightarrow \mathcal{I}_Z(r) \rightarrow \mathcal{O}(r) \rightarrow \mathcal{O}_Z \rightarrow 0$. Since $r \leq 0$, the only possibility is $r = 0$ and $\deg Z = 1$. In conclusion, if E doesn't split, we have: $0 \rightarrow \mathcal{O} \rightarrow E \rightarrow \mathcal{I}_P(-1) \rightarrow 0$, where P is a point. Such bundles do exist.

Remark 5. *One can show the following: let E be a stable, rank two vector bundle on \mathbb{P}^2 , with $c_1(E) = -1$.*

If $h^1(E) = 0$ then there exists an exact sequence: $0 \rightarrow \mathcal{O} \rightarrow E(1) \rightarrow \mathcal{I}_Z(1) \rightarrow 0$, where Z is a set of three non collinear points. We have $c_2(E) = 3$.

If $h^1(E(1)) = 0$ then there exists an exact sequence: $0 \rightarrow \mathcal{O} \rightarrow E(2) \rightarrow \mathcal{I}_Z(3) \rightarrow 0$, where Z is a set of six points not lying on a conic. We have $c_2(E) = 4$.

Let us recover Theorem 1.

Lemma 6. (1) *Let \mathcal{F} be a stable, normalized, rank two vector bundle on \mathbb{P}^3 . Then $h^1(\mathcal{F}(-1)) \neq 0$.*

(2) *Moreover if $c_1(\mathcal{F}) = -1$, we have $h^1(\mathcal{F}).h^1(\mathcal{F}(1)) \neq 0$.*

Proof. (1) Let $H \subset \mathbb{P}^3$ be a general plane and consider the exact sequence $0 \rightarrow \mathcal{F}(-2) \rightarrow \mathcal{F}(-1) \rightarrow \mathcal{F}_H(-1) \rightarrow 0$. Assume $h^1(\mathcal{F}(-1)) = 0$. By Barth's restriction theorem ([1]) $h^0(\mathcal{F}_H(-1)) = 0$. It follows that $h^1(\mathcal{F}(-2)) = 0$ and then $h^1(\mathcal{F}(-m)) = 0, m \geq 1$. Now we have $h^2(\mathcal{F}(-2)) = h^1(\mathcal{F}(-c_1 - 2)) = 0$. This implies $h^1(\mathcal{F}_H(-1)) = 0$ and by Theorem 2, \mathcal{F}_H splits. This implies that \mathcal{F} also splits (see [9]), a contradiction. Hence $h^1(\mathcal{F}(-1)) \neq 0$.

(2) Assume $h^1(\mathcal{F}) = 0$. By stability we have $h^3(\mathcal{F}) = h^0(\mathcal{F}(-3)) = 0$. It follows that $\chi(\mathcal{F}) = h^2(\mathcal{F}) \geq 0$. By Riemann-Roch we get $1 - 3c_2/2 \geq 0$. This is impossible since $c_2 > 0$ and c_2 is even.

Assume $h^1(\mathcal{F}(1)) = 0$. We have $h^3(\mathcal{F}(1)) = h^0(\mathcal{F}(-4)) = 0$. It follows that $\chi(\mathcal{F}(1)) \geq 0$. By Riemann-Roch this yields: $5 - 5c_2/2 \geq 0$. Since c_2 is even and $c_2 > 0$, it follows that

$c_2 = 2$. Stable rank two vector bundles on \mathbb{P}^3 with $c_1 = -1, c_2 = 2$ have been classified ([7]) and they all have $h^1(\mathcal{F}(1)) = 1$. \square

Lemma 7. *Let \mathcal{F} be a non-stable, normalized, rank two vector bundle on \mathbb{P}^3 . If $h^1(\mathcal{F}(-1)) = 0$, then \mathcal{F} splits.*

Proof. Since \mathcal{F} is not stable we have an exact sequence: $0 \rightarrow \mathcal{O} \rightarrow \mathcal{F}(r) \rightarrow \mathcal{I}_C(2r + c_1) \rightarrow 0$ (*), where $r \leq 0$ and where C is either empty or a l.c.i. curve with $\omega_C(4 - 2r - c_1) \simeq \mathcal{O}_C$ (**). Assume $h^1(\mathcal{F}(-1)) = 0$ and C non empty. Twisting by $\mathcal{O}(-r - 1)$ and taking cohomology, we get $h^1(\mathcal{I}_C(r - 1 + c_1)) = 0$. Since $r - 1 + c_1 < 0$ this implies $h^0(\mathcal{O}_C(r - 1 + c_1)) = 0$. It follows from (**) that $h^0(\omega_C(-r + 3)) = 0$. Now consider the exact sequence: $0 \rightarrow \mathcal{I}_C(r - 2 + c_1) \rightarrow \mathcal{I}_C(r - 1 + c_1) \rightarrow \mathcal{I}_{C \cap H}(r - 1 + c_1) \rightarrow 0$, where H is a general plane. If $h^2(\mathcal{I}_C(r - 2 + c_1)) = 0$, then $h^1(\mathcal{I}_{C \cap H}(r - 1 + c_1)) = 0$. Restricting (*) to H and twisting by $-r - 1$, we get $h^1(\mathcal{F}_H(-1)) = 0$. By Theorem 2, \mathcal{F}_H splits, hence \mathcal{F} also splits, which contradicts the minimality of the twist r (C should be empty). So $h^2(\mathcal{I}_C(r - 2 + c_1)) = h^1(\mathcal{O}_C(r - 2 + c_1)) \neq 0$. By Serre duality on C : $h^1(\mathcal{O}_C(r - 2 + c_1)) = h^0(\omega_C(-r + 2 - c_1)) \neq 0$. But this contradicts $h^0(\omega_C(-r + 3)) = 0$. We conclude that C is empty and that \mathcal{F} splits. \square

Lemma 8. *Let \mathcal{F} be a non stable rank two vector bundle on \mathbb{P}^3 , with Chern classes $c_1 = -1, c_2$. If $h^1(\mathcal{F}) = 0$ or $h^1(\mathcal{F}(1)) = 0$, then \mathcal{F} splits.*

Proof. Since \mathcal{F} is not stable we have an exact sequence: $0 \rightarrow \mathcal{O} \rightarrow \mathcal{F}(r) \rightarrow \mathcal{I}_C(2r - 1) \rightarrow 0$, with $r \leq 0$. Twisting by $\mathcal{O}(m)$ and taking cohomology, we see that $h^0(\mathcal{F}(m+r)) = h^0(\mathcal{O}(m))$ as long as $m \leq -2r + 1$ (since then $h^0(\mathcal{I}_C(m + 2r - 1)) = 0$). Twisting by \mathcal{O}_H , H a general plane, we get $0 \rightarrow \mathcal{O}_H \rightarrow \mathcal{F}_H(r) \rightarrow \mathcal{I}_{C \cap H}(2r - 1) \rightarrow 0$. Arguing as above we get that $h^0(\mathcal{F}_H(m+r)) = h^0(\mathcal{O}_H(m))$ if $m \leq -2r + 1$. We conclude that the exact sequence $0 \rightarrow \mathcal{F}(k-1) \rightarrow \mathcal{F}(k) \rightarrow \mathcal{F}_H(k) \rightarrow 0$ is exact on H^0 if $k \leq -r + 1$. In particular we have $0 \rightarrow H^1(\mathcal{F}(k-1)) \rightarrow H^1(\mathcal{F}(k))$, if $k \leq -r + 1$. If $h^1(\mathcal{F}(t_0)) = 0$ with $t_0 \leq -r + 1$, then $h^1(\mathcal{F}(m)) = 0$ for $m \leq t_0$. So if $h^1(\mathcal{F}).h^1(\mathcal{F}(1)) = 0$, then $h^1(\mathcal{F}(-1)) = 0$. Since $h^2(\mathcal{F}(-2)) = h^1(\mathcal{F}(-1))$, we get $h^1(\mathcal{F}_H(-1)) = 0$. By Theorem 2 we conclude that \mathcal{F}_H splits, hence \mathcal{F} also splits. \square

Putting every thing together we get:

Proposition 9. *Theorem 2 implies Theorem 1.*

Remark 10. *The original proof in [3] has been worked out in the framework of sub-canonical space curves.*

Let us conclude this section with a last remark:

Proposition 11. *Let E be an indecomposable rank two vector bundle on \mathbb{P}^2 . Then the module $H_*^1(E)$ is connected (i.e. if $h^1(E(t)) \neq 0$ and $h^1(E(m)) \neq 0$ with $t < m$, then $h^1(E(k)) \neq 0$ for $t < k < m$).*

According to Theorem 2 this is equivalent to the following: (a) if $h^1(E(-t)) = 0$ for some $t \geq 2$, then $h^1(E(-m)) = 0, \forall m \geq t$, and (b) if $h^1(E(t)) = 0$ for some $t \geq 0$, then $h^1(E(m)) = 0, \forall m \geq t$.

Proof. (1) First assume E stable. Using the exact sequence $0 \rightarrow E(-t-1) \rightarrow E(-t) \rightarrow E_L(-t) \rightarrow 0$ ($L \subset \mathbb{P}^2$ a general line) and the fact that $h^0(E_L(-t)) = 0$ if $t \geq 2$ (because $E_L \simeq \mathcal{O}_L \oplus \mathcal{O}_L(c_1)$, by stability), condition (a) follows immediately.

Now (b) follows from (a) by duality, indeed $h^1(E(t)) = h^1(E(-t-c_1-3))$ and $t+3+c_1 \geq 2$.

(2) Assume E non stable. Then we have an exact sequence $0 \rightarrow \mathcal{O} \rightarrow E(r) \rightarrow \mathcal{I}_Z(2r+c_1) \rightarrow 0$, with $r \leq 0$, $Z \subset \mathbb{P}^2$ zero-dimensional. If $h^1(E(t)) = 0, t \geq 0$, then $h^1(\mathcal{I}_Z(2r+c_1+t)) = 0$. Since Z is zero-dimensional we have $h^1(\mathcal{I}_Z(k)) = 0, \forall k \geq 2r+c_1+t$, hence $h^1(E(m)) = 0, \forall m \geq t$. This proves (b). Now (a) follows by duality: by assumption $0 = h^1(E(-t)) = h^1(E(t-c_1-3))$. Since $t \geq 2, t-c_1-3 \geq 0$, except if $c_1 = 0, t = 2$ but this case cannot occur since $h^1(E(-1)) = h^1(E(-2)) \neq 0$ by Theorem 2. So if $c_1 = 0$, we may assume $t \geq 3$. \square

Remark 12. (i) *This improves Castelnuovo-Mumford's lemma at least for the vanishing part.*

(ii) *It can be shown that the H^1 -module of an indecomposable rank two vector bundle on \mathbb{P}^3 is connected, but the proof is much more difficult, see [2].*

3. RANK TWO VECTOR BUNDLES ON \mathbb{P}^2 WITH $h^1(E(-1)) \leq 4$.

In this section we will investigate bundles with $h^1(E(-1)) =: u$ small, say $u \leq 4$. Let us start with a useful remark:

Remark 13. *Assume E indecomposable, r as usual and consider $0 \rightarrow \mathcal{O} \rightarrow E(r) \rightarrow \mathcal{I}_Z(2r+c_1) \rightarrow 0$, where $Z \subset \mathbb{P}^2$, is zero-dimensional. Let $0 \rightarrow \mathcal{L}_1 \rightarrow \mathcal{L}_0 \rightarrow \mathcal{I}_Z \rightarrow 0$ be the minimal free resolution of \mathcal{I}_Z . Then we can lift the morphism $\mathcal{L}_0(2r+c_1) \rightarrow \mathcal{I}_Z(2r+c_1)$ to a morphism $\mathcal{L}_0(2r+c_1) \rightarrow E(r)$ and then get (after a twist) an exact sequence:*

$$(3) \quad 0 \rightarrow \mathcal{L}_1(r+c_1) \rightarrow \mathcal{O}(-r) \oplus \mathcal{L}_0(r+c_1) \rightarrow E \rightarrow 0$$

This gives the minimal free resolution of $H_^0(E)$. Now by dualizing and taking into account that $E^* = E(-c_1)$ we get:*

$$(4) \quad 0 \rightarrow E \rightarrow \mathcal{O}(r+c_1) \oplus \mathcal{L}_0^*(-r) \rightarrow \mathcal{L}_1^*(-r) \rightarrow 0$$

Taking cohomology we get the beginning of the minimal free resolution of the $S := k[x, y, z]$ module $H_^1(E)$:*

$$0 \rightarrow H_*^0(E) \rightarrow S(r+c_1) \oplus L_0^*(-r) \rightarrow L_1^*(-r) \rightarrow H_*^1(E) \rightarrow 0$$

Then combining with (3) we get the whole resolution. By the way we notice a curious fact: $rk(S(r + c_1) \oplus L_0^*(-r)) = rk(L_1^*(-r)) + 2$. So for a finite length graded module M , to be the H^1 -module of a rank two vector bundle on \mathbb{P}^2 , the number of relations among its generators must be the number of generators plus two. In fact this is not only necessary but also sufficient (see [10] for details).

Lemma 14. *Let E be a normalized rank two vector bundle on \mathbb{P}^2 . Assume E indecomposable, with $h^1(E(-1)) =: u$. Let r be the minimal twist of E having a section. If E is not stable, then $E(r)$ has a section vanishing on a zero-dimensional subscheme, Z , with $\deg(Z) = u$.*

Proof. We have an exact sequence $0 \rightarrow \mathcal{O} \rightarrow E(r) \rightarrow \mathcal{I}_Z(2r + c_1) \rightarrow 0$, with $r \leq 0$ since E is not stable. Twisting by $\mathcal{O}(-r + 1)$ and taking cohomology we get: $h^1(E(-1)) = h^1(\mathcal{I}_Z(r + c - 1 - 1)) = h^0(\mathcal{O}_Z)$, because $r + c_1 - 1 < 0$ (notice that $-r - 1 \geq -1$, hence $h^2(\mathcal{O}(-r - 1)) = 0$). It follows that $\deg(Z) = u$. \square

Remark 15. (1) *In view of this lemma and on Remark 13 if we know all the possible minimal free resolutions of u points we get all possible resolutions of $H_*^0(E)$. Observe that the minimal free resolution of $H_*^0(E)$ determines the whole cohomology of E . Indeed if we know $h^0(E(k)), \forall k \in \mathbb{Z}$, then by duality we know $h^2(E(k)), \forall k \in \mathbb{Z}$. Knowing $h^0(E(k)), h^2(E(k))$, we get $h^1(E(k))$ by Riemann-Roch.*

(2) *If E is non stable, indecomposable, then $h^0(E(r)) = 1$, hence $Z = (s)_0$ is uniquely defined. So we can define a map, γ , from the set of non stable bundles with $h^1(E(-1)) = u$ to $Hilb^u(\mathbb{P}^2)$, by $\gamma(E) = Z$.*

Lemma 16. *Let E be a stable, normalized, rank two vector bundle on \mathbb{P}^2 . We have $u := h^1(E(-1)) = c_2$.*

Proof. Since $h^0(E(-1)) = 0 = h^2(E(-1)) = h^0(E(-c - 1 - 2))$, we have $\chi(E(-1)) = -h^1(E(-1))$. By Riemann-Roch $\chi(E(-1)) = -c_2$ and the result follows. \square

Remark 17. *At this point the classification, or better the description, of rank two vector bundles E with $h^1(E(-1)) = u$ can be split into two parts:*

(1) *for non stable bundles: it is enough to determine all the minimal free resolutions of l.c.i., zero-dimensional subschemes of degree u .*

(2) *classification of stables vector bundles of Chern classes $-1 \leq c_1 \leq 0$ and $c_2 = u$. In particular we want to know the least twist having a section.*

Observe that the set of non stable bundles with $h^1(E(-1)) = u$ is some kind of counterpart to the moduli space $\mathcal{M}(c_1, c_2)$ ($c_2 = u$) in the stable case.

Let us start with non stable bundles. To make things manageable we will assume $u \leq 4$.

Lemma 18. *Let $Z \subset \mathbb{P}^2$ be a closed subscheme of codimension two, with $\deg(Z) = u \leq 5$. There are ten possible minimal free resolutions for the ideal of Z , namely:*

(a) Z is contained in a line, in this case Z is a complete intersection $(1, u)$

(b1) $u = 3$ and Z is not contained in a line, in this case:

$$0 \rightarrow 2.\mathcal{O}(-3) \rightarrow 3.\mathcal{O}(-2) \rightarrow \mathcal{I}_Z \rightarrow 0.$$

(b2) $u = 4$, $h^0(\mathcal{I}_Z(1)) = 0$, but Z has a subscheme of length three contained in a line. In this case: $0 \rightarrow \mathcal{O}(-3) \oplus \mathcal{O}(-4) \rightarrow 2.\mathcal{O}(-2) \oplus \mathcal{O}(-3) \rightarrow \mathcal{I}_Z \rightarrow 0$

(b3) $u = 4$ and Z is a complete intersection $(2, 2)$.

(b4) $u = 5$, $h^0(\mathcal{I}_Z(1)) = 0$ but Z has a subscheme of length 4 contained in a line. In this case: $0 \rightarrow \mathcal{O}(-3) \oplus \mathcal{O}(-5) \rightarrow 2.\mathcal{O}(-2) \oplus \mathcal{O}(-4) \rightarrow \mathcal{I}_Z \rightarrow 0$

(b5) $u = 5$, $h^0(\mathcal{I}_Z(2)) = 1$. In this case:

$$0 \rightarrow 2.\mathcal{O}(-4) \rightarrow 2.\mathcal{O}(-3) \oplus \mathcal{O}(-2) \rightarrow \mathcal{I}_Z \rightarrow 0$$

Proof. Well known. □

As explained before this gives us all the possible resolutions (hence all the possible cohomologies) of non stable, indecomposable bundles with $h^1(E(-1)) \leq 4$. We need $u = 5$ for the stable case:

Proposition 19. *Let E be a stable, normalized, rank two vector bundle on \mathbb{P}^2 , with $h^1(E(-1)) = u \leq 4$. As usual let r denote the minimal twist of E having a section. Then $r = 1$ or $r = 2, c_1 = -1, u = 4$. Moreover:*

(1) *If $c_1 = 0$ we have $u \geq 2$ and $E(1)$ has a section vanishing on a subscheme of degree $u + 1$ which is not contained in a line.*

(2) *If $c_1 = -1$ and $r = 1$, we have $u \geq 1$ and $E(1)$ has a section vanishing on a subscheme of length u . If $r = 2, u = 4$, then $E(2)$ has a section vanishing on a degree 6 subscheme, Z , with $h^0(\mathcal{I}_Z(2)) = 0$.*

Proof. In any case $h^2(E(1)) = h^0(E(-c_1 - 4)) = 0$ by stability. Since $\chi(E(1)) = 6 - c_2$ if $c_1 = 0$ (resp. $4 - c_2$ if $c_1 = -1$) and since $c_2 = u$ (Lemma 16), we get $\chi(E(1)) > 0$, except if $c_1 = -1, u = 4$. In this case we have $h^0(E(1)) = h^1(E(1))$. Assume $h^0(E(1)) = 0$. Since $\chi(E(2)) = 9 - c_2$, we have $h^0(E(2)) > 0$ and an exact sequence $0 \rightarrow \mathcal{O} \rightarrow E(2) \rightarrow \mathcal{I}_Z(3) \rightarrow 0$, where $\deg(Z) = c_2(E(2)) = 6$. We have $h^0(E(1)) = 0 = h^0(\mathcal{I}_Z(2))$. This proves the first claim.

(1) Assume $c_1 = 0$. We have $0 \rightarrow \mathcal{O} \rightarrow E(1) \rightarrow \mathcal{I}_Z(2) \rightarrow 0$. By stability $h^0(\mathcal{I}_Z(1)) = 0$. In particular $\deg Z = c_2 + 1 \geq 3$, i.e. $u = c_2 \geq 2$.

(2) Assume now $c_1 = -1$. Since $c_2(E(1)) = c_2 = u$, if $r = 1$, we have $0 \rightarrow \mathcal{O} \rightarrow E(1) \rightarrow \mathcal{I}_Z(1) \rightarrow 0$, with Z of degree u . If $u = 4$ and $h^0(E(1)) = 0$, a section of $E(2)$ vanishes along Z of degree 6 with $h^0(E(1)) = 0 = h^0(\mathcal{I}_Z(2))$. □

Remark 20. (1) *By Serre's construction for any $k \leq 2$ and any locally complete intersection, zero-dimensional subscheme $Z \subset \mathbb{P}^2$ there exists a rank two vector bundle, F ,*

with $c_1(F) = k$, and an exact sequence $0 \rightarrow \mathcal{O} \rightarrow F \rightarrow \mathcal{I}_Z(k) \rightarrow 0$. If $k \leq 0$ this is the least twist of F having a section and F is not stable. In particular all the bundles we have considered in Proposition 19 do really exist !

(2) We have the list of all possible resolutions for $H_*^0(E)$, where E is a normalized bundle with $h^1(E(-1)) \leq 4$. Indeed the only case not covered by Lemma is when $r = 2$, but a subscheme, Z , of degree 6, not on a conic has a resolution like: $0 \rightarrow 3\mathcal{O}(-4) \rightarrow 4\mathcal{O}(-3) \rightarrow \mathcal{I}_Z \rightarrow 0$.

(2) We observe that if $u = 1$ we always have that $E(r)$ has a section vanishing at one point. More precisely:

Corollary 21. *Let E be a normalized, indecomposable, rank two vector bundle on \mathbb{P}^2 . Let r denote the minimal twist of E having a section.*

(1) *The following are equivalent:*

- (i) $h^1(E(m)) \leq 1, \forall m \in \mathbb{Z}$
- (ii) $h^1(E(-1)) = 1$
- (iii) $E(r)$ has a section vanishing at one point
- (iv) there is an exact sequence:

$$0 \rightarrow \mathcal{O}(-b-1) \rightarrow \mathcal{O}(-a) \oplus 2\mathcal{O}(-b) \rightarrow E \rightarrow 0$$

with $a \leq b$ (in particular $a = r, b = -r - c_1 + 1$).

(2) *A bundle like in (1) is stable if and only if $a = b$, if and only if $E = \Omega(1)$.*

Proof. (i) \Leftrightarrow (ii), since E is indecomposable, this follows from Theorem 2.

(ii) \Rightarrow (iii): If E is non stable this follows from Lemma 14. If E is stable this follows from Proposition 19. More precisely we have $c_2 = 1$ (Lemma 16) and $r = 1, c_1 = -1$ (Proposition 19).

(iii) \Rightarrow (iv): This follows from Remark 13.

(iv) \Rightarrow (i): Since $a \leq b, r = a$, hence a section of $E(a)$ will vanish in codimension two. Since $c_1 = -r - b + 1$, we get $b = -r - c_1 + 1, c_1(E(a)) = a - b + 1$. We get a commutative diagram:

$$\begin{array}{ccccccc}
 & & & 0 & & 0 & \\
 & & & \downarrow & & \downarrow & \\
 & & & \mathcal{O} & = & \mathcal{O} & \\
 & & & \downarrow & & \downarrow & \\
 0 & \rightarrow & \mathcal{O}(a-b-1) & \rightarrow & \mathcal{O} \oplus 2\mathcal{O}(-b+a) & \rightarrow & E(a) \rightarrow 0 \\
 & & \parallel & & \downarrow & & \downarrow \\
 0 & \rightarrow & \mathcal{O}(a-b-1) & \rightarrow & 2\mathcal{O}(-b+a) & \rightarrow & \mathcal{I}_Z(a-b+1) \rightarrow 0 \\
 & & & & \downarrow & & \downarrow \\
 & & & & 0 & & 0
 \end{array}$$

So we get $0 \rightarrow \mathcal{O}(-2) \rightarrow 2\mathcal{O}(-1) \rightarrow \mathcal{I}_Z \rightarrow 0$ and we conclude that Z is a point p . Since $h^1(\mathcal{I}_p(m))$ is 0 if $m \geq 0$ and 1 if $m < 0$, we conclude that $h^1(E(k)) \leq 1, \forall k \in \mathbb{Z}$.

(2) We have already seen ((ii) \Rightarrow (iii)) that E is stable if and only if $c_2 = r = 1, c_1 = -1$.

Hence we have $0 \rightarrow \mathcal{O}(-1) \rightarrow 3\mathcal{O} \rightarrow E(1) \rightarrow 0$. It follows that $E(1) = T(-1) = \Omega(2)$. On the other hand if $r = a = b = -r - c_1 + 1$, then $c_1 = -1$ and $r = 1$, in particular E is stable. \square

Remark 22. *This result is known in the context of logarithmic bundles, see [4], [8]*

In the same vein we have:

Corollary 23. *Let E be a normalized, indecomposable, rank two vector bundle on \mathbb{P}^2 . Let r denote the minimal twist of E having a section. The following are equivalent:*

- (i) $h^1(E(-1)) = 2$
- (ii) $E(r)$ has a section vanishing along a subscheme of degree two, or E is stable with $c_1 = 0, c_2 = 2, r = 1$ and $E(1)$ has a section vanishing along a subscheme of degree three not contained in a line.
- (iii) there is an exact sequence:

$$0 \rightarrow \mathcal{O}(-b-2) \rightarrow \mathcal{O}(-b-1) \oplus \mathcal{O}(-b) \oplus \mathcal{O}(-a) \rightarrow E \rightarrow 0$$

with $a \leq b$ (in particular $a = r, b = -r - c_1 + 1$), or:

$$0 \rightarrow 2\mathcal{O}(-2) \rightarrow 4\mathcal{O}(-1) \rightarrow E \rightarrow 0.$$

Proof. It is similar to the previous one, so we omit it. \square

REFERENCES

- [1] Barth, W.: *Some properties of stable rank-2 vector bundles on \mathbb{P}^n* , Math. Ann., **226**, 125-150 (1977)
- [2] Buraggina, A.: *The intermediate cohomology modules of a rank two vector bundle on \mathbb{P}^3 are not split*, Comm. Alg., **25**, 999-1008 (1997)
- [3] Chiantini, L.-Valabrega, P.: *Subcanonical curves and complete intersections in projective 3-space*, Ann. Mat. Pura Appl., **138**, 309-330 (1984)
- [4] Dimca, A.-Sticlaru, G.: *Nearly free divisors and rational cuspidal curves*, arXiv:1505.0066v3 [math.AG] 2 Jun 2015 (2015)
- [5] Ein, L.: *An analogue of Max Noether's theorem*, Duke Math. J., **52**, 689-706 (1985)
- [6] Ellia, Ph.: *Quasi complete intersections and global Tjurina number of plane curves*, preprint arXiv (2019)
- [7] Hartshorne, R.-Sols, I.: *Stable rank two vector bundles on \mathbb{P}^3 with $c_1 = -1; c_2 = 2$* , J. reine u. angewandte Math., **325**, 145-152 (1981)
- [8] Marchesi, S.-Vallès, J.: *Nearly free curves and arrangements: a vector bundle point of view*, arXiv:1712.0486v2 [math.AG] 11 Oct 2018 (2018)
- [9] Okonek, Ch.-Schneider, M.-Spindler H.: *Vector bundles on complex projective spaces*, Progress in Mathematics **3**, Birkhäuser (1980)
- [10] Rao, P.: *A note on cohomology modules of rank two vector bundles*, Journal of Algebra, **86**, 23-34 (1984)

DIPARTIMENTO DI MATEMATICA E INFORMATICA, UNIVERSITÀ DEGLI STUDI DI FERRARA, VIA MACHIAVELLI 30, 44121 FERRARA, ITALY.

E-mail address: phe@unife.it